

Contents

Preface to the second edition	ix
To the student ... please read this first	xi
Choosing books for further work	xv
Acknowledgements	xix

EM ESSENTIAL MATHS

EM1 Numbers and symbols	2
1 Positive and negative numbers: adding and subtracting	4
2 Positive and negative numbers: multiplying and dividing	6
3 Combining addition, subtraction, multiplication and division	10
4 Using a calculator	14
5 Introducing letters and symbols	20
6 Working with symbols: adding, subtracting, multiplying, dividing	27
EM2 Simplifying expressions	33
1 Equivalent fractions and cancelling	34
2 Adding and subtracting fractions	40
3 Multiplying and dividing fractions	46
4 Putting it all together – and a note on percentages	54
5 Expanding brackets	57
6 Factorising	64
7 Powers	70
8 Powers of products and quotients	80
9 Fractional powers	85
EM3 Solving problems	93
1 How equations arise	94
2 Solutions to equations: recognising and guessing	97
3 A method for solving equations	101
4 Formulating and solving equations	108
5 Rearranging equations and substituting	111
6 Inequalities	123
EM4 Modelling using straight lines	133
1 Introducing straight lines	134
2 Interpretation of a and b	140
3 Linear equations for modelling	146
4 Pairs of linear equations	150

MM MORE MATHS

MM1	Some special equations	160
1	Introducing logs	161
2	Solving equations when the unknown is a power	168
3	Quadratic equations	174
4	The expert equation solver	181
MM2	Modelling using curves	188
1	Functions	189
2	Quadratic curves	192
3	Some common functions	203
4	Variations on standard curves	208
5	When do these curves arise?	212
6	Curve sketching in general	217
MM3	Rates of change	220
1	The gradient of a curve	221
2	More differentiation	229
3	Interpreting derivatives	234
4	Maximising and minimising	242

DD DESCRIBING DATA

DD1	Pictures of data	256
1	Frequencies and histograms	257
2	Stem and leaf diagrams	267
3	Pictures of one categorical variable: bar charts and pie charts	271
4	More graphs for numerical data	277
5	Categorical data that come in pairs	281
6	Lies, more lies and statistics?	286
DD2	Summarising data	291
1	The centre of a set of data	292
2	The spread of a set of data	298
3	Σ and a short-cut for variance	304
4	Quartiles	309
5	Technology to the rescue!	311
6	Constructing indices	314
7	Grouped data	318
8	Describing data: your toolkit	324

P PROBABILITY

P1	Measuring uncertainty	328
1	Introduction to probability	329

2	Combining events: AND and OR	344
3	Conditional probability	352
4	Calculating joint or AND probabilities	360
5	More complicated probabilities – made simple!	368
P2	Numerical outcomes	375
1	Introducing random variables	376
2	Expectation: the long run average	384
3	The variance of a random variable	391
4	The binomial distribution	398
5	The Poisson distribution	415
P3	Continuous numerical outcomes	426
1	Probability density functions (pdfs)	427
2	The exponential distribution	440
3	Introducing the normal distribution	449
4	Calculating normal probabilities	460
5	The normal approximation to the binomial distribution	471

S STATISTICS

S1	Estimation	483
1	Samples	484
2	Estimating from a sample	490
3	How good is an estimator?	496
4	Interval estimates	505
5	Beware, variance unknown!	516
S2	Testing hypotheses	525
1	Introduction to testing	526
2	The structure of a test	533
3	Tests for the mean	539
4	Rejection regions: another approach to testing	555
S3	Correlation and regression	563
1	Data that come in pairs	564
2	Fitting a straight line to the data	574
3	The linear regression model	585
4	Extending the linear regression model: the multiple linear regression model	602
S4	Forecasting	611
1	Displaying time series	612
2	Introduction to forecasting	616
3	Coping with trend and seasonal effects	627
4	Forecasting series with a trend and seasonal effect	644
S5	Comparing two populations	648
1	The difference between two means	649
2	Difference between two means: small samples	657

3 Paired samples	663
S6 Categorical data	672
1 Proportions	674
2 More than two categories	680
3 Contingency tables	689
Statistics formulae	699

BM BUSINESS MODELLING

BM1 Linear programming models	705
1 A small linear programming problem	707
2 More complicated linear programming models	715
BM2 Planning projects	725
1 Drawing a project planning network	726
2 Scheduling activities	733
BM3 Models for inventory control	742
1 An inventory model	743
2 Making the model more realistic	749
BM4 Time and money	757
1 Calculating interest	758
2 Present values	767
3 Series of payments	775
BM5 Decision making	789
1 Introducing decision making	790
2 Payoff tables	793
3 Decision-making criteria	797
4 Decision trees	804
BM6 Controlling quality	820
1 Control charts for the mean	821
2 <i>R</i> -charts and <i>p</i> -charts	830
BM7 Simulating reality	839
1 Why simulate?	840
2 A simple simulation	841
3 Generating random inputs	847
4 Simulating from other probability distributions	853
Statistical tables	861
Index	874

EM1

Numbers and symbols

The concept of number is the obvious distinction between beast and man. Thanks to number, the cry becomes song, noises acquire rhythm, the spring is transformed into a dance, force becomes dynamic, and outlines figures.

Joseph Marie de Maistre, French author

Contexts

What is this chapter about?

This chapter explains why numbers are important, both in society and in business. It reminds you about negative numbers and explains how to add, subtract, multiply and divide them and how to use brackets. It explains how to make rough estimates before using a calculator and how to round numbers. It goes on to describe how symbols can be used to represent general relationships between quantities and shows how symbols can be treated in a very similar way to numbers. It introduces the idea of a model and explains what a spreadsheet is.

Why is it useful?

Numbers are used everywhere to describe and measure, to allocate resources and to plan ahead. So the basic numeracy skills in this chapter are an essential skill of modern life. Most professional jobs these days require you to use a spreadsheet and a calculator and in Business, Management and Finance, you will frequently have to manipulate numbers and understand simple mathematical expressions which involve symbols, for instance to represent how budgets are allocated to different departments or calculate an interest payment. And any quantitative subject you study either in education or for professional examinations will require the work in this chapter.

Where does it fit in?

The first part of the chapter describes the basic operations of arithmetic, representing numbers and using a calculator, whereas the second part, using symbols instead of numbers, is really the very beginning of the subject of algebra. Your memories of school algebra might put you off but please bear with us – we show you that symbols can be used in exactly the same way as numbers and we start from the very beginning.



What do I need to know?

We assume only that you know how to add, subtract, multiply and divide positive numbers and zero and that you can cope with numbers like 5.123 in which the fractional part is given using decimal places. Most readers will have covered the work in this part of the book when they were younger. We realise that this may have been a long time ago and that not everyone got on with maths at school, so the explanations are thorough and completely self-contained.

Objectives

After your work on this chapter you should be able to:

- add, subtract, multiply and divide positive, negative and zero numbers;
- combine the operations above using brackets;
- make rough estimates before using a calculator;
- understand how to round to so many decimal places or significant figures and how to use scientific notation;
- use symbols to represent the relationships between quantities;
- understand the difference between constants and variables;
- evaluate an expression using particular values for the variables;
- understand the concept of a model;
- understand what a spreadsheet is;
- perform basic operations to simplify expressions containing symbols.

Imagine a world without numbers. Accurate measurement would be impossible. Physical phenomena like air temperature, medical diagnostics such as blood pressure or economic statistics like the inflation rate and unemployment figures would be impossible to quantify. We would be left saying vaguely that ‘prices seem to have gone up’, or ‘tomorrow will be warmer’, but further analysis and comparison would be impossible.

Our society allocates resources – raw materials, labour and property – almost entirely by ascribing monetary values to them. Without numbers your prospective career would not exist – no accounts, no economic models, no sales figures.

So we need numbers. They enable us to describe exactly how our world is now, to allocate resources and to plan into the future.

In management, you will have to represent quantities numerically, calculate them, analyse relationships between them and communicate your findings to clients and colleagues. This will often require you to use symbols to represent the quantities of interest. For instance, a general rule for calculating the amount received at the end of a year, on an investment of £ A at an annual interest rate of $r\%$ is

$$A \times \left(1 + \frac{r}{100}\right)$$

(Don't worry if you don't understand this now – we will see how it is obtained later.)

In the first four sections of this chapter we will only use *numbers* but in the remaining sections we will use symbols as well. The ways of dealing with symbols are just the same as those for numbers – hardly surprising when you think that the symbols are standing in for numerical values anyway.

1 Positive and negative numbers: adding and subtracting

Here is the first diagnostic 'test box'. Take a minute to try it. If you can answer correctly, with no difficulty whatsoever, you could move directly on to Section 2.

Test box 1

Can you solve these?


$$2 + (-5) =? \quad -3 - (-4) =? \quad 3 - (-2) + (-5) =?$$

Solutions: -3, 1, 0.

Negative numbers

You are already familiar with numbers, and with decimal places. Numbers like 3, 1000, 10.24, and 3.1427 are all *positive* numbers; that is, they are *greater* than 0. For many applications (for example, temperatures in Celsius below freezing, profit and loss or credits and debits in a bank account) and as a tool for many mathematical skills it is useful to be able to talk about *negative* or *minus* numbers. These are numbers like -5, -7.74, -1000 or -1.0. We will often enclose them in brackets like (-5) or (-7.74) to show that the minus sign belongs to that number.

All these numbers, positive, negative and zero, can be represented on a line. Part of it is shown below. The dots at each end show that the line could continue forever in both directions.



Adding and subtracting negative numbers

We all know that $4 + 5 = 9$, or $8 - 3 = 5$, but it is less obvious how to add and subtract negative numbers.

Any sum can be represented by a traveller on a journey along the line above. He starts at one point on it and from there can go forwards (to the right) to higher numbers or backwards (to the left) to lower numbers. Adding and subtracting represent forward and backward progress respectively. For instance, for $(-5) + 3$, the traveller starts at -5 on the line and then moves 3 units forwards to arrive at -2, so $(-5) + 3 = (-2)$. For $(-4) - 1$ she starts at -4 and moves one step backwards to (-5) so $(-4) - 1 = (-5)$.

Check this

Use the line to convince yourself that $(-3) + 2 = (-1)$, that $(-3.7) + 1.7 = (-2)$ and that $(-1) - 3 = -4$.

When we need to add or subtract a *negative* number the direction must be reversed because the number is negative. For example, to calculate $3 + (-2)$, he starts at 3 on the line, would move forward for the addition but does the reverse because we are adding a negative number. So he moves 2 units backwards from 3 to arrive at 1 and we conclude that $3 + (-2) = 1$. In a similar way, to calculate $3 - (-2)$ he starts at 3, would move backwards for the subtraction but does the reverse because of the -2 and so moves 2 units forwards to 5.

Check this

Convince yourself that $(-3) + (-2) = (-5)$ and that $(-1) - (-2) = 1$.

Notice that as $3 + (-2)$ results in moving 2 units backwards it gives exactly the same result as $3 - 2$ and that as $3 - (-2)$ results in moving 2 units forwards it is the same as $3 + 2$. This is because 'forwards' and 'backwards' are the reverse of each other. For $3 - (-2)$ the effect is one of a double negative, the phrase 'I am *not not* going' or the opposite of an opposite.

So, plus a minus number is the same as minus a positive number (opposite signs), whereas minus a negative number is the same as plus a positive number (same signs) as shown below.

Adding and subtracting negative numbers

OPPOSITE SIGNS

$+ (- \text{number})$ or $- (+ \text{number})$ gives a $-$

SAME SIGNS

$- (- \text{number})$ or $+ (+ \text{number})$ gives a $+$

When evaluating sums it is usually easiest to rewrite them in terms of positive numbers only as we have done in the following examples.

Check these

$$\begin{aligned} 3 + (-5) \\ = 3 - 5 = -2. \end{aligned}$$

$$\begin{aligned} 1 - (-4) \\ = 1 + 4 = 5. \end{aligned}$$

$$(-3) - 7 = -10.$$

$$12.42 - (-3.1) \\ = 12.42 + 3.1 = 15.52.$$

When more than two numbers appear in a sum we work in the same way. For instance $3.2 - (-3) + (-2) - 1 = 3.2 + 3 - 2 - 1 = 3.2$.

Check this

$$\begin{array}{r} -6.7 + (-7) - (-0.1) + 2.1 \\ = -6.7 - 7 + 0.1 + 2.1 = -11.5 \end{array}$$

2 Positive and negative numbers: multiplying and dividing

Test box 2

Can you do these?

$$\begin{array}{l} (-3) \times 2 \quad (-4) \times (-0.5) \quad -2 \times 10 \quad 8 \div (-2) \\ -6 \div (-3) \quad -4 \times 12 \times (-2) \end{array}$$

Solutions: (row-wise) $-6, 2, -20, -4, 2, 96$.

Multiplying negative numbers

Multiplication means 'times' so 2×3 is really 'two threes' or $3 + 3$, 4×3 is 'four threes' $3 + 3 + 3 + 3$ and so on. The order of multiplication does not matter so 2×3 is the same as $3 \times 2 = 2 + 2 + 2$, and 4×3 is the same as $3 \times 4 = 4 + 4 + 4$ and so on.

The number line in Section 1 can help us to work out how to multiply negative numbers.

The multiplication 2×3 is $3 + 3$, so our traveller starts at 0 and travels forward 3, and then forward 3 again to reach 6. By similar reasoning $2 \times (-3) = (-3) + (-3)$, so he starts at 0 and then travels *backwards* 3 steps and then backwards 3 steps again to reach -6 , so we have that $2 \times (-3) = -6$. If the negative number comes first, we can reverse the order of multiplication. For instance, to calculate $(-4) \times 2$ we regard this as $2 \times (-4)$ and work out $(-4) + (-4) = (-8)$.

The big problem comes when we want to multiply two negative numbers together, let's say $(-5) \times (-2)$. How can our traveller do backward steps of 2, *minus* 5 times? Or backward steps of 5, *minus* 2 times? Here we do need a leap of faith. The convention is that *a negative times a negative is a positive*. This rule was adopted because everything then falls into a pattern for later work.

The key results for multiplying positive and negative numbers are:

Multiplying numbers

of the same sign gives a +
of a different sign gives a –

that is,

$$+ \times + = +$$

$$+ \times - = -$$

$$- \times + = -$$

$$- \times - = +$$

Of course, to multiply more than two numbers together we just multiply the first two, then multiply the result by the third number and so on. A property of multiplication is that we will obtain the same result regardless of the order in which we multiply the numbers. For instance,

$$\begin{aligned} 2 \times 3 \times 4 \times 5 \\ = 6 \times 4 \times 5 \\ = 24 \times 5 = 120 \end{aligned}$$

or we could have said

$$\begin{aligned} 2 \times 3 \times 4 \times 5 \\ = 2 \times 12 \times 5 \\ = 2 \times 60 = 120 \end{aligned}$$

The result of multiplying two or more numbers together is called the *product* of those numbers. For instance 6 is the product of 2 and 3 and 100 is the product of 5 and 20.

Check this

What is the product of 6 and 30? Of 2 and –5 and 20?

Answers: 180 and –200.

Dividing negative numbers

The rules for division have to comply with those for multiplication. For example, because $4 \times 5 = 20$, 20 divided by 4 must be 5 and 20 divided by 5 must be 4. So the rules we have already met for multiplication using negative numbers dictate the rules for division. For instance, as $4 \times (-5) = -20$, we can deduce that $(-20) \div (-5) = 4$ and that $(-20) \div 4 = -5$.

Dividing numbers

of the same sign gives a +
of a different sign gives a –

that is,

$$\begin{aligned} + \div + &= + \\ + \div - &= - \\ - \div + &= - \\ - \div - &= + \end{aligned}$$

Notice that the rules for the sign of a division using negative numbers are just the same as the rules for multiplication; that is, dividing numbers with *different* signs gives a negative result, whereas dividing those of the *same* sign gives a positive result.

Check these

$$10 \div (-5) = (-2)$$

$$-100 \div 20 = -5$$

$$-28 \div (-7) = 4.$$



Dividing by 0

It is *not* possible to divide by 0. How can a quantity be split into 0 parts?

Alternatives to the \div sign

You may remember that division can be written in several ways. The division $8 \div 2$ can also be written $\frac{8}{2}$ or $8/2$ and the result is called the *quotient*. Writing a division in this way is particularly useful when a whole expression needs to be divided by another whole expression. For instance, the division of $80 - 20$ by $5 + 10$ can be written

$$\frac{80 - 20}{5 + 10}$$

Be especially careful about the exact length of the quotient line. For instance

$$1 + \frac{5 + 3}{2}$$

(which equals 5) is crucially different from

$$\frac{1 + 5 + 3}{2}$$

(which equals 4.5).

- Evaluate the following:
 - $6 + (-3) - (-4)$
 - $(-3) + (-7) - 11 - (-8)$
 - $0 - (-3) - (-1) - (-4)$
 - $-4 + 3 - (-2)$
 - $-5 - (-2) + (-4)$
- Evaluate the following:
 - $6 \times (-3)$
 - -4×5
 - $(-8) \times (-8)$
 - $6 \times (-1)$
 - $(-5) \times (-10)$
 - $(-60) \times (-2)$
 - $(-2) \times (-60)$
 - What is the product of 2 and 20?
- Evaluate the following:
 - $10 \div (-2)$
 - $60 \div (-15)$
 - $-12 \div 6$
 - $-12 \div (-6)$
 - $\frac{12}{6}$
 - $\frac{12}{-6}$
 - $\frac{-12}{6}$
 - $\frac{-12}{-6}$
- Evaluate the following:
 - $(-4) \times (-3) \times (-5)$
 - $(-4) \times 3 \times (-5)$
 - $(-1) \times (-1) \times (-1) \times (-1) \times (-1) \times (-1)$
 - $(-1) \times (-1) \times (-1) \times (-1) \times (-1) \times (-1) \times (-1)$
 - $-1 \times -5 \times -10$
 - $-1 \times -5 \times 10$

Solutions:

- a. 7 b. -13 c. 8 d. 1 e. -7
- a. -18 b. -20 c. 64 d. -6 e. 50 f. 120 g. 120 h. 40
- a. -5 b. -4 c. -2 d. 2 e. 2 f. -2 g. -2 h. 2
- a. -60 b. 60 c. 1 d. -1 e. -50 f. 50

- Calculate the following:
 - $(-20) - (-10) - (-5)$
 - $0 - 20 - (-10) + (-5)$
 - $(-1) - (-1) - 1 + (-1)$
- Evaluate:
 - $10 \times (-5)$
 - $(-2) \times 6$
 - $(-5) \times (-10) \times (-3)$
 - $\frac{-200}{10}$
 - $\frac{200}{-10}$
 - $-2 \times -2 \times -2 \times -2 \times -2$
 - $-2 \times -2 \times -2 \times -2 \times -2 \times -2 \times -2$
 - $\frac{-60}{-3}$
 - $\frac{-40}{8}$
 - $\frac{40}{-8}$
 - $\frac{-40}{-8}$

3 Combining addition, subtraction, multiplication and division

Test box 3

$$4 \times 8 - 6 \div 2 \quad 3 \times (50 - 10) \quad \left(\frac{40 - 20}{10 - 5} \right) \times 3$$

$$((75 \div 25) + 2) \times 5$$

Solutions: 29, 120, 12, 25.

The order of operations

Calculate $3 - 2 \times 4$.

It may look simple enough, but there is a snag. There are two ways to proceed and they each give a different answer. You might have reasoned:

(i) subtract first to give $3 - 2 = 1$ and then calculate $1 \times 4 = 4$ so the answer is 4

or you might have said

(ii) multiply first, so $2 \times 4 = 8$, and $3 - 8 = -5$.

Which one did you do?

The answer depends on the order in which the calculations are performed – whether to subtract or multiply first – and so longer expressions may have many more than two alternative answers. This situation is unsatisfactory so we need some sort of rule which tells us in which order to perform the operations.

The accepted rule is to *multiply and divide first*, performing calculations from left to right and *then add and subtract*, also from left to right. So for the example above we should multiply first, so the second answer, $3 - 2 \times 4 = 3 - 8 = -5$, is correct.

Let's try $6 \times 2 \div 4 + 1$. The rule tells us to multiply and divide first, but there is both a multiplication and a division here, so we must work from left to right. The multiplication occurs before the division, so we multiply 6×2 first to give

$$12 \div 4 + 1,$$

then divide giving

$$3 + 1,$$

and finally do the addition,

$$= 4.$$

Check these

Remember to perform calculations from left to right if there is more than one multiplication/division or more than one addition/subtraction.

$$\begin{aligned}
 & -2 + 7 \times 8 \div 2 \times 2 \\
 = & -2 + 56 \div 2 \times 2 \\
 = & -2 + 28 \times 2 \\
 = & -2 + 56 = 54 \\
 \\
 & 50 - 32 \div 16 \times 2 \\
 = & 50 - 2 \times 2 \\
 = & 50 - 4 = 46
 \end{aligned}$$

Introducing brackets

Suppose we need to multiply $2 + 3$ by $4 - 2$. We *cannot* write this as

$$2 + 3 \times 4 - 2$$

because applying the order of operations rule would give $2 + 12 - 2 = 12$. To show that the numbers must be processed in a different order we use *brackets*.

When part of an expression must be evaluated first it must be enclosed in brackets. So the multiplication of $2 + 3$ by $4 - 2$ must be written $(2 + 3) \times (4 - 2)$. This indicates that we must first calculate $2 + 3$, and then calculate $4 - 2$, and finally multiply the results $5 \times 2 = 10$.

As evaluating the expressions in brackets takes priority over anything else, we can extend the rule for the order of operations to *brackets*, multiply and divide, add and subtract.

For example, to evaluate $(9 - 2) \times 10 - (2 \times 3)$ we work out the brackets first to give

$$7 \times 10 - 6$$

then multiply to give

$$70 - 6$$

and finally subtract to obtain

$$= 64.$$

The order of operations in evaluating arithmetic expressions is

Brackets

Multiply and Divide (from left to right)

Add and Subtract (from left to right)

When multiplying by a bracket it is usual to omit the multiplication sign. So $(2 + 3) \times (4 - 2)$ is written $(2 + 3)(4 - 2)$ and $2 \times (4 + 7)$ is written $2(4 + 7)$.

Check these

$$\begin{aligned}
 & 6 \quad (4 - 6) \quad (4 + 2) \quad + \quad 3 \\
 = & 6 \quad \times(-2) \quad \times 6 \quad + \quad 3 \\
 = & \quad (-12) \quad \times 6 \quad + \quad 3 \\
 = & \quad \quad (-72) \quad + \quad 3 \quad = -69 \\
 & 36 \quad \div \quad (4 \times 3) \quad - \quad 5 \\
 = & 36 \quad \div \quad 12 \quad - \quad 5 \\
 = & \quad 3 \quad - \quad 5 \quad = -2
 \end{aligned}$$

Quotients can be written using brackets, for instance,

$$\frac{80 - 20}{10 - 5}$$

can be written $(80 - 20) \div (10 - 5)$ because everything *above* the line of a quotient is divided by everything *below* the line.

Sometimes more than one layer of brackets is necessary. Do not be put off by this. You will find that you have to work out the *inside* brackets first and then proceed outwards. For instance

$$(6(1 + 4)) \div 10 = (6 \times 5) \div 10 = 30 \div 10 = 3.$$

Authors and lecturers are sometimes helpful and use different symbols for different 'layers' of brackets. For instance $\{[(2 + 3) + 5] \times 7\}$. As you will not always encounter this we have often used only one symbol in our work.

Check this

$$\begin{aligned}
 & 10 \quad \times \quad (2 \quad + \quad (6 \div 3) \quad \times \quad 4) \\
 = & 10 \quad \times \quad (2 \quad + \quad 2 \quad \times \quad 4) \\
 = & 10 \quad \times \quad (2 \quad + \quad 8) \\
 = & 10 \quad \times \quad 10 \quad = 100
 \end{aligned}$$

In practice, brackets are often used to clarify expressions when they are not strictly essential.

WORK CARD 3

1. Evaluate the following:

a. $(20 - 5) \times (4 - 2)$

b. $2 + (10 \div 5) \times 3$

c. $2 \times (10 \div 5) \times 2$

d. $2 \times 10 \div (5 \times 2)$

e. $\frac{10 + 20}{4 - 2}$

f. $2 \times 2 \times (27 \div 3) + (1 - 20)$

g. $(4 \times 2 \times 2) + (5 \times (-1))$



2. Evaluate the following:

a. $1 + 3 \times (4 + (8 \div 2))$

b. $((50 \div 25) \times 8 \div (7 - 3)) \times 3$

c. $12 - (4(8 \times (6 - 4)) - 5)$

d. $\frac{18 \times (2 - 3 \times 4)}{(4 + 14)}$

e. $\left(\frac{18 \div 3}{(4 \times 3) - 36}\right) \times 4$

f. $(-10) \times \left(\left(\frac{100}{25} \times 2\right) + (60 \div 20)\right) + 1$

g. $1 + \left(\frac{2 \times (2 + 4 \times 5)}{(2 \times 11) \div (22 \div 2)}\right)$

Solutions:

1. a. 30 b. 8 c. 8 d. 2 e. 15 f. 17 g. 11

2. a. 25 b. 12 c. -47 d. -10 e. -1 f. -109 g. 23

1. Evaluate:

a. $(40 \div 2) + (3 \times 4)$

b. $-5 + (-3) \times 2 + 1$

c. $6 \div (3 \times 2) + 4$

d. $3 \times 3 \times (6 \div 2) + 3$

e. $3 \times 3 \times 6 \div 2 + 3$

f. $\frac{3+6}{5-2}$

g. $\frac{10-2}{3+1} \div (-2)$

2. Evaluate:

a. $(21 \div 7) + (50 \div (5 \times 2) + 1)$

b. $\frac{77 \div 11}{108 \div (-3) \times 4}$

c. $\left(\frac{40}{6+2} \left(3 + \frac{5 - (21 \div 7)}{6-4} \times \frac{48}{16}\right)\right) + 5$

d. $\left(20\,000 \div \left((1000 - (2 \times 5 \times 50)) \times \left(\frac{100}{50}\right)\right)\right) - 2$

e. $\frac{30 + 3 \times 3}{10 + 3} + 5 \times \frac{10 + 100}{11}$

4 Using a calculator

Test box 4

1. Use a calculator to evaluate:

$$\frac{500}{1.2(20 + 34)} \text{ to 6 decimal places.}$$

2. Express the following to 3 decimal places:

$$1.9755 \quad 10,002.9999 \quad 209.452 \quad 12.73 \quad 0.000123456$$

3. Express the numbers in question 2 to 3 significant figures.

4. Express the following in scientific notation:

$$12,000,000 \quad 0.00001254$$

5. By estimating roughly, do you think that

$$\frac{515 \times 6.1}{200} = 7.1$$

is correct?

Solutions:

1. 7.716049 2. 1.976 10,003.000 209.452 12.730 0.000

3. 1.98 10,000 209 12.7 0.000123 4. 1.2×10^7 1.254×10^{-5}

5. Use rough estimates: $500 \times 6 = 3000$, divided by 200 gives 15, the answer doesn't look good.

Which sort of calculator?

You will need a fairly basic calculator with the usual $+$, $-$, \times , \div and also simple functions such as $\frac{1}{x}$, \sqrt{x} , x^2 , \log and \ln (natural logarithm), 10^x , e^x , x^y . A memory would also be helpful.

The order of operations on a calculator

Like any other invaluable tool a calculator is only as good as its operator – the modern adage, 'garbage in garbage out' is extremely relevant here. The calculator will only produce the right answer if you supply the numbers and operations in the correct order – which may not be the order in which they are written.

Try to evaluate the following expressions using your calculator:

$$(20 + 30) \div 3$$

$$20 + (30 \div 3)$$

The answers you should have are 16.666666 and 30. To calculate the first expression you need to enter $20 + 30$ on your calculator and then divide by 3. For the second one you need

to evaluate the bracket $30 \div 3$ first and *then* add 20. We should add that some calculators do provide bracket functions, which we suggest that you use with care.

Show your working

Think about the errors you make when you word-process, or type or write! These are usually apparent when you read through later. It is just as easy to press the wrong key on a calculator, but most machines won't display all your inputs. It is therefore a good idea to write out your intermediate workings and your train of thought for a problem and not just the final value the calculator gives you. By doing this mistakes are easier to spot, your work is easier to follow – for a colleague or for yourself later – and last but not least, if your answer is wrong but your method is right you will still get most of the available marks in an exam.



Rough estimates

Always keep in mind the *real* problem you are solving.

Be critical of the answer your calculator gives you. Look out for percentage decreases that are over 100, negative probabilities, or fractions when you expected whole numbers.

Other errors may be less obvious and so it is a good idea to perform a rough mental calculation to get an idea of the magnitude of the solution.

To illustrate this, consider the following scenarios – and their solutions. Which ones seem reasonable and which don't? *Don't* use a calculator.



Check these

1. A university has 6782 students, about half of which can be expected to visit a campus catering outlet on any given term-time day. Lunches are £2–3 and coffee and a snack is about £1. The total takings for a day over all outlets is £6142. Does this seem reasonable?
2. At a bank there is a single queueing system and five cashiers. On average, during peak hours a customer arrives every 32 seconds. The situation is modelled mathematically (such models are called queueing models) to assess the effects of increasing or reducing the number of cashiers on duty. The final result from the model shows that on average 76% of customers would have to wait longer than five minutes if there were 4 cashiers, 56% if there were 5 cashiers and 66% if there were 6 cashiers.
3. I earn \$8.42 an hour and worked $96\frac{1}{2}$ hours last month. My payslip for the month says \$585.19. Does this seem reasonable?
4. The interest I will earn on £6179 invested at a rate of 5% for the first £5000 and 7.2% on the remainder over a year is given by

$$5000 \times \frac{5}{100} + 1179 \times \frac{7.2}{100}$$

Using a calculator I obtain £334.88. Does this seem right?

Solutions:

1. Working in thousands, roughly 4000 students will visit an outlet, and if an equal number have coffee or lunch they will spend an average of about £1.75, making total revenue for the day about £7000. The result is about right.

2. The results are suspicious here as more cashiers should bring down the percentage of people who have to wait longer than 5 minutes, whereas here the figure for 6 cashiers is greater than for 5. Maybe the modelling procedure is inappropriate, or else a calculation is erroneous.
3. Approximately \$8 for roughly 100 hours should give me \$800, so something is wrong. In fact this wage is for 69.5 hours!
4. Yes, it seems reasonable. Say that the average rate of interest is about 6% and that the sum invested is about £6000. We would expect the interest to be about

$$6000 \times \frac{6}{100} = \text{£}360$$

Rounding: decimal places and significant figures

When you use a calculator or computer to perform a calculation the machine will only display a certain number of digits. The exact answer may need many more digits or even an infinite number of them. For instance, when we divide 5 by 17 our calculator shows 0.294117647.

Numbers can be rounded to a particular number of *decimal places* (*d.p.*) or a particular number of *significant figures* (*sig. fig.*).

The convention for rounding to a particular number of decimal places (*d.p.*) is that when the first digit to be *excluded* is between 5 and 9, we round up, and when it is between 0 and 4 we round down. So, for instance, 3.625 expressed to 2 decimal places rounds up to 3.63, and 3.624999 rounds down to 3.62. Remember to include 0s where appropriate. For instance, 3.634999 to 4 decimal places is 3.6350, and to 5 decimal places is 3.63500.

A second way of representing numbers approximately is to write them to a particular number of *significant figures* (*sig. fig.*). The left-most digit of a number is the most significant as it is the digit that represents the greatest value, the next from left-most is the second most significant and so on. So in the number 672.34 the '6' represents hundreds and is therefore the most significant figure, the '7' represents the number of tens and is the second most significant figure and so on.

To write a number to, say, 3 significant figures we use the three left-most digits, rounding the final one if necessary. As when rounding to so many decimal places, if the first *discarded* digit is 5 or more we round up and if it is between 0 and 4 we round down. For example, 6248.500052 to 3 sig. fig. is 6250, to 2 sig. fig. is 6200 and to 8 sig. fig. is 6248.5001.

Check this

What is 7,254,600 to 3 sig. fig., to 4 sig. fig?
 What is 0.00652445 to 3 sig. fig., to 5 sig. fig.?

Solutions: 7,250,000, 7,255,000, 0.00652, 0.0065245.

Notice from the last example that any zeros after the decimal point but before the first non-zero digit do not count as 'significant'.

Most calculators display numbers to 8 or 10 significant figures.

In practice, the accuracy with which we need to record results depends on the application. You do not see newspaper headlines reporting that the inflation rate is

3.42534213%: it is usually given to just one decimal place; that is, 3.4%. A chemist analysing a substance may have measuring equipment that is only accurate to 0.001g so it is pointless to record the result to 6 decimal places. A statistical model that forecasts the percentage dividend payable by a company for the next 10 years to 32 decimal places is itself only an approximation, so it is meaningless to report the forecasts to more than perhaps 1 or 2 decimal places.

So, common sense must prevail when deciding how many decimal places or significant figures to give in the *answer* to a problem. However, it can be dangerous to round too much *during* your calculations. Consider the following example.



Past records show that 8892 2cm tacks were manufactured at a cost of £13.16. A management accountant needs to estimate the price at which the factory should sell a batch of 20,000 tacks and reasons as follows.

Cost of manufacturing 8892 tacks is £13.16.

$$\text{Cost per tack is } \frac{13.16}{8892} = \text{£}0.001 \text{ rounded to 3 d.p.}$$

A batch of 20,000 tacks therefore costs $20,000 \times 0.001 = \text{£}20$.

She concludes that if the factory sells a batch for £25 it will make a profit.

This is obviously an extreme example and we hope that you can see the source of error here. The accurate calculations are Cost per tack is

$$\frac{13.16}{8892}$$

so a batch of 20,000 tacks costs

$$20,000 \times \frac{13.16}{8892} = \text{£}29.60$$

so the factory would make a *loss* if it sold a batch for £25.

Computers and calculators perform all their calculations to a particular accuracy. When you write out calculations by hand you may round your intermediate values to a different number of decimal places and so your results may differ slightly from those of the machine.

Scientific notation

You may have noticed that when your calculator is faced with a very large or a very small number it resorts to another notation called *scientific notation*. For instance, calculate $123,456,789 \times 1234$ on your machine.

Our calculator display shows

$$1.523456776^{11}$$

The small raised 11 means 'multiplied by 10 to the power of 11'.

We will cover powers more thoroughly in Chapter EM2, but for now it is enough to know that 10 to the power of 11 is written 10^{11} and means 10 multiplied by itself 11 times, or

$$10 \times 10 \times 10 \times 10 \times 10 \times 10 \times 10 \times 10 \times 10 \times 10 \times 10.$$

So the number above can be written $1.523456776 \times 10^{11}$.

To multiply a number by 10 you may recall that we just move the decimal point one place to the right; for example, $1.523456776 \times 10 = 15.23456776$. So to multiply by 10, eleven times, we move the decimal point 11 places to the right. The number represented above is therefore 152345677600.0. Remember that this result is not likely to be exact as the calculator only displays a certain number of significant figures.

Very small numbers can be represented using negative powers of 10. Again, these are described in Chapter EM2. Multiplying by 10^{-5} is the same as dividing by 10^5 or 100,000. We can do this by moving the decimal point 5 places *to the left*. So if the diameter of an atomic particle is 0.00000000000000000014 mm it is much easier to write it as 1.4×10^{-20} , where multiplying a number by 10^{-20} equates to moving the decimal point 20 places to the left.

1. Use a calculator to evaluate the following:

$$\frac{24.15}{5(150 + 11)} \quad \frac{42 - 20.04}{366 \times 24} \quad \frac{120 \div 15}{85 + 35}$$

2. *Without* using a calculator, decide which of the following answers are most likely:

a. $\frac{312.42 \times 7.54}{21} = 112.17366$ or 1.1217 or 0.01122

b. 112 articles are purchased at £5.42 and a further 62 articles at £2.42. The total bill is £757.08, £1050.12 or £342.01?

c. I purchase 300 euros at an exchange rate of 1.5 euro to £1. This costs me £200.00, £350.00 or £450.00?

d. In a traffic census at an accident black spot, 938 cars are seen to pass in an hour. Of these, 300 are going at over 50 kilometres per hour (kph) and 638 are travelling more slowly. The average speed of these cars is calculated to be 36.4 kph, 51 kph or 16 kph?

3. Write the following to 3 decimal places:

$$1.9755 \quad 10,002.9752 \quad 209.452 \quad 12.73 \quad 0.00012456$$

4. Express the numbers in question 3 to 3 significant figures.

5. Express the following in scientific notation

$$12,000,000 \quad 0.00001254 \quad 9.999999999 \quad 999.999$$

6. Write the following numbers out in full

$$2.678 \times 10^6 \quad 4.1 \times 10^{-9} \quad 1 \times 10^0$$

Solutions:

1. 0.03 0.0025 0.06̇ (remember the 6̇ means that the digit 6 repeats forever – it's called 6 *recurring*).
2. Use your calculator to check a., b. and c. In d. the only reasonable answer is 36.4.



WORK CARD 4

3. 1.976 10,002.975 209.452 12.730 0.000
4. 1.98 10,000 209 12.7 0.000125
5. 1.2×10^7 1.254×10^{-5} 9.999999999×10^0 9.99999×10^2
6. 2,678,000 0.0000000041 1

ASSESSMENT 4

1. Evaluate the following using a calculator:

$$\frac{312 \times 101}{42.7 - 21.5} \quad 432.2 - (543.2 - 10.17)/2.5$$

2. Say which of the following results seem sensible and explain why. Do *not* use a calculator except to check your answer.

a. A leisure aircraft is owned by a Flying Group. The Group estimate that maintenance costs are likely to be £1000 next year. The main running cost is fuel which currently costs £0.7 per litre, but will be £0.95 per litre next year. Last year the aircraft did 9000 km. Average fuel consumption is 2.9 km per litre. If the 10 members of the Group are to share costs equally, the Treasurer suggests that a suitable monthly cost per member which ensures that maintenance and fuel costs are covered is £32.90, £44.12 or £12.10.

b. $\frac{111 + (20.5 \times 10.7)}{302} = 10.939, 1.0939 \text{ or } 0.10939$

c. I give myself a budget of \$100 a day for my 8-day holiday. At the start of my holiday I have £600 which I change into dollars at a rate of \$1.57 = £1. During the holiday I do not exceed my budget. At the end of the holiday I calculate that I should have \$40 or \$142 or \$10 left. (Ignore commission charges.)

3. Express the following to 4 decimal places

$$13.66666 \quad -3.156 \quad 200,000.00001 \quad 156.99999 \quad 55.12345$$

4. Express the numbers in question 3 to 2 significant figures.

5. Write the following in scientific notation

$$3,000,051.0 \quad 0.0000009142 \quad -102.01 \quad 4.14$$

6. Write the following numbers out in full

$$3.42 \times 10^8 \quad 1.004 \times 10^{-6} \quad 9.99 \times 10^3$$

5 Introducing letters and symbols

Test box 5

The total cost of a holiday for four friends comprises the cost of four return flights at \$ f each and a charge of \$70 per day for a rented cottage. Write down an expression for the cost of the holiday for each person, if they go for d days.

Evaluate $\frac{2st}{s+1}$ when $s = 2$ and $t = 6$.

Evaluate $3a + b$, when $a = 4$ and $b = 5$.

Simplify $\frac{a}{b} + 2 - 5a - 3\frac{a}{b} + a$

What is meant by 'modelling'? Have you ever used a spreadsheet?

Solutions:

$$\frac{4f + 70d}{4} \text{ or } f + \frac{70d}{4}, \quad 8, \quad 17, \quad -2\frac{a}{b} + 2 - 4a$$

See the sub-sections on 'models' and 'using a spreadsheet' below.

Why use letters and symbols?

Suppose we have £200 and wish to purchase some euros. The exchange rate is £1 = 1.5 euros and for simplicity we will ignore any commission charge. Using a calculator we can work out that the £200 will buy $200 \times 1.5 = 300$ euros.

That's fine – but exchange rates fluctuate and not everyone wants to change exactly £200. It would be much more useful to develop a *general relationship* between the number of pounds, the exchange rate and the number of euros.

To obtain the figure of 300 euros we *multiplied* the number of pounds by the exchange rate so the relationship we seek is

$$\text{euros} = \text{pounds} \times \text{rate}$$

This relationship is valid for any number of pounds and any exchange rate. So to calculate, for instance, how many euros we would obtain when the exchange rate is 1.9 and we have £300 to spend we write out the relationship again:

$$\text{euros} = \text{pounds} \times \text{rate}$$

but *substitute* 300 instead of 'pounds' and 1.9 instead of 'rate' to give

$$\text{euros} = 300 \times 1.9$$

and so euros = 570. We would obtain 570 euros.

It is usual to use a single letter to represent each of the entities in a relationship. Letters like x and y are often used but sometimes we use 'meaningful' letters like e for euros, p for pounds and r for rate, in which case our relationship would be written

$$e = p \times r$$

However, it is common to omit the multiplication sign adjacent to symbols – just as we omit multiplication signs next to brackets – and so we could write this as

$$e = pr$$

A relationship like $e = pr$ is an *equation* (because it contains an = sign), or we could say that the *formula* for the number of euros, e , is pr .

Once we have an equation or formula it can be used in different ways to solve a variety of problems. For instance, suppose we know that we received 960 euros at the Bureau de Change in exchange for £800, but we can't remember what the exchange rate was. Substituting $e = 960$ and $p = 800$ into the equation above gives

$$960 = 800r$$

We now need to *solve* the equation for the unknown quantity r . We will see how to do this in Chapter EM3, but for now we will tell you that the solution is

$$r = \frac{960}{800} = 1.2$$

so the exchange rate was £1 = 1.2 euros.

This currency example is relatively uncomplicated but it has demonstrated that symbols are useful

- (i) to represent the general relationship between the quantities of interest; and so
- (ii) to enable calculation of one quantity from the others.

Using symbols to represent relationships

We now concentrate on how to turn information on the quantities of interest into expressions involving symbols. There is no magic way of doing this but if you find it hard we suggest that you break the task into two steps as follows.

1. Read through the information you have been given, but as you come across each quantity, assign it a symbol. Make a written note of these.
2. Read through again, but as you read 'translate' each fact you are given into symbols.

We include some examples with commentary. Remember that multiplication signs are usually omitted next to symbols.

Check these

1. It costs £2000 (the fixed cost) to set up a production run in a factory, and then a further £5 (the variable cost per unit) for each item manufactured. Write down an expression for the total production cost.

Solution:

On a first reading we realise that we need a symbol for the number of items manufactured (say, n) and a symbol for the total production cost (say, C).

During a second read-through we make the following jottings, which culminate in the desired expression for total production cost.

Fixed cost	2000
n items at a cost of £5 per item $5 \times n$	<u>$5n$</u>
Total cost $C =$	$2000 + 5n$

2. A restaurant has two menus, a tourist menu at £8 and a gourmet menu at £15. Write down a formula for the cost of the food for a party in which t customers have the tourist menu and g customers choose the gourmet menu.

Solution:

The symbols t and g have already been chosen for us here, but we adopt C for the total cost. Reading through we would write down something like

Tourist menu £8	t customers	8t
Gourmet £15	g customers	<u>15g</u>
Total cost is $C =$		$8t + 15g$

3. I want to organise a group of friends to hire a boat on the river for the day. The basic hire cost is £60, but we must also pay for fuel which costs £5 an hour. Write down an expression for the cost per person, C . (This might be of interest to establish how many friends I need to ask to keep the cost per person down to a particular amount.)

Solution:

Suppose I ask $n - 1$ friends so there are n of us altogether, and hire a boat for h hours. The total cost will be $60 + 5h$, so the cost per person will be

$$C = \frac{60 + 5h}{n}$$

Constants and variables

In many applications some amounts will be fixed – like the cost of fuel in 3 or the price of the meal in 2. These fixed amounts are called *constants*. They are usually numbers, although a symbol which represents a particular value, like $\pi = 3.14159$ (from the formula for the circumference of a circle, $2\pi r$, where r is the radius) is also a constant.

On the other hand, symbols that represent quantities that can change, such as the number of people n , or the number of hours, h in 3, are called *variables*.

Evaluating expressions

Once you have an expression you will often need to calculate its value when the variables in it take particular values.

For instance, suppose the amount of tax payable on a salary of S at a tax rate of $t\%$ is given by

$$\frac{(S - 3000)t}{100}$$

If Paul earns a salary of £10,000 and the tax rate is 30% then he must pay

$$\frac{(10,000 - 3000) \times 30}{100} = \text{£}2100$$

All we have done is replace the symbols in the formula with the values we are interested in. This is called *substitution*.

Check this

The amount of interest I receive on an investment of £ P at $r\%$ interest over one year less a management charge of £10 + 0.01 P is

$$\frac{P(r - 1)}{100} - 10$$

Calculate the amount of interest I would receive in the following cases

- On an investment of £10,000 when the interest rate is 5%.
- On an investment of £500 when the interest rate is 10%.

Solution:

$$\text{a. } \frac{10,000 \times (5 - 1)}{100} - 10 = \text{£}390$$

$$\text{b. } \frac{500 \times (10 - 1)}{100} - 10 = \text{£}35$$

Models

Much of the work you will do for your degree and beyond will require you to build financial, accounting and economic models. By a *model* we usually mean one or more equations which represent the real-life situation.

The advantage of a model is that it allows us to answer ‘what if ...’ type questions, without changing the real system. We illustrate this with a simple model for a clothing manufacturer’s business.

Suppose it costs \$10,000 to design a particular item of ladies clothing, and the unit production cost is \$15. We might model the profit as

$$\text{Profit} = \text{price} \times \text{number sold} - (10,000 + 15 \times \text{number manufactured})$$

The model enables the manufacturer to calculate the profit for a variety of values of the unknown variables (price, number sold and number manufactured) so that he may choose the values for price and number manufactured that give the highest profit. (He is not likely to have any control on the number sold.)

A model does not pretend to be exact. It will often be based on (and only as good as) a series of assumptions that the modeller makes (and should state), but the hope is that the model is a reasonable approximation to the real situation.

A model may be as complex or as simple as we like. For instance, the model above makes the assumption that the number sold is unaffected by the price. It would probably be more realistic to assume that as the price increased the number of sales decreased. This could be built into the model by introducing another relationship, for instance

$$\text{number sold} = 1000 - (10 \times \text{price})$$

and the two relationships could be used together to find the best price and number to manufacture.



Using a spreadsheet

For those of you who are *not* familiar with spreadsheets, they are a type of computer software that comprises a grid of numerical quantities and the relationship between them. They have become a useful tool for building models in accountancy, business, finance and elsewhere.

The rows of a spreadsheet are numbered 1, 2, 3, ..., etc. and columns are labelled A, B, C, ..., etc. so that each cell of the grid is uniquely identified by a letter and a number like A1, B23 or E2.

Into a cell of a spreadsheet the user can either enter a number – in which case the cell always takes that value – or an expression giving the relationship between the current cell and the others. For instance, if I would like the total of cells E12 and E13 to appear in cell E14 I would enter the expression

$$= E12 + E13$$

into cell E14. The exact format of this expression may differ depending on the spreadsheet package, but the principle is always the same.

Spreadsheets are programmed so that when a value in a cell is changed, all the values elsewhere in the spreadsheet, which are influenced by that cell's value, will be changed automatically. Thus, 'what if ...' questions can be answered by trial and error.

The spreadsheet screen can show two modes – one which shows the symbolic expressions in each cell and the other which gives the numerical values implied by these.

A simple spreadsheet which calculates student marks is shown below.

	A	B	C	D	E	F
1	student	c w 1 %	c w 2 %	overall c w	exam %	overall %
2	Catherine	58	66	$= (B2+C2)/2$	77	$= 0.2*D2+0.8*E2$
3	Sylvia	82	70	$= (B3+C3)/2$	45	$= 0.2*D3+0.8*E3$
4	Malcolm	55	45	$= (B4+C4)/2$	76	$= 0.2*D4+0.8*E4$
5	Dennis	78	70	$= (B5+C5)/2$	60	$= 0.2*D5+0.8*E5$
6	Veronica	90	82	$= (B6+C6)/2$	74	$= 0.2*D6+0.8*E6$
7	Gillian	60	62	$= (B7+C7)/2$	58	$= 0.2*D7+0.8*E7$
8						
9					average	$= (F2+F3+F4+F5+F6+F7)/6$

20% of the marks on a Maths course can be gained from coursework and 80% from the exam. The two pieces of coursework carry equal weight and are each marked out of 100. A spreadsheet which calculates and displays the overall coursework mark and the overall marks of each student, and calculates the class's average mark is shown above in formula mode. Notice that the expressions for the overall coursework marks require brackets.

The same spreadsheet in 'values' mode is shown below.

student	c w 1 %	c w 2 %	overall c w	exam %	overall %
Catherine	58	66	62	77	74
Sylvia	82	70	76	45	51.2
Malcolm	55	45	50	76	70.8
Dennis	78	70	74	60	62.8
Veronica	90	82	86	74	76.4
Gillian	60	62	61	58	58.6
				average	65.63

In this book we will make use of a spreadsheet program, Microsoft Excel, particularly for *Statistics* and *Business Modelling*.

WORK CARD 5

- Write down expressions for the following. Start by naming the variables you need.
 - The cost of an m metre length of a roll of carpet when the width of the roll of carpet is 4 m and the cost per square metre is £12.
 - The cost of a group's meal at a pizzeria, when pizzas are £4 each and bottles of wine £7. No other food or drink is available.
 - The amount a gardener earns in a week when he charges £5 per hour for manual work like weeding but £8 per hour for skilled gardening work.
 - The net amount given or charged to me in one month by my bank when they give me 5% interest on my average balance over the month, but charge me 30p per debit transaction.
 - The cost of petrol for a journey of L miles when my average fuel consumption is 30 miles per gallon and the cost of fuel is £ p per litre. Assume there are 4.54 litres in a gallon.
- Salaries in a computer consultancy firm are calculated according to the following formula.

$$S = 8000 + 300(A - 20) + 1000Y$$

where A is the age of the employee in years, and Y is the number of years of experience they have in computing.

What will an employee's salary be if she

- Has just graduated at 21 with no work experience?
 - Left school at 16 and has worked in computing for 4 years until today?
 - Is now 60, but took up Computing work at the age of 52 for the first time?
- A company has lent a sum of £ P . It is to be repaid at $r\%$ interest in one year's time. The company calculates that the prevailing market interest rate is $i\%$, so the present value to the company of the repayment is



$$V = \frac{P(1 + \frac{r}{100})}{1 + \frac{i}{100}}$$

Find the present value in each of the following cases:

- The loan is £10,000 at 8% interest and the market interest rate is 10%.
- The loan is £15,000 at 12% interest and the market interest rate is 10%.
- The loan is £20,000 at 5% interest and the market interest rate is 5%.

Solutions:

- $48m$
 - $4p + 7w$
 - $5m + 8s$
 - $0.05A - 0.3D$
 - $\frac{L}{30} \times 4.54p$ (L is length of journey in miles)
- 8300
 - 12,000
 - 28,000
- £9818.18
 - £15,272.72
 - £20,000.

- Write down an expression for the floor area of a two-room flat when the first room has one and a half times the width and twice the breadth as the second. Start by naming the variables you are going to use.
- I want to buy some Botswana pulas and the exchange rate is 8.25 BWP = £1 if I buy them abroad using a credit card or 8.5 BWP = £1 for a cash transaction at a Bureau de Change. The credit card company will charge me 2.3% commission on a credit card transaction whereas a Bureau de Change charges 2% commission plus a fee of £3.

Write down an expression for the cost (including commission) of buying a particular number of Botswana pulas (a) by credit card and (b) from a Bureau de Change.

 - Evaluate each of these expressions for 700 BWP, 800 BWP and 1000 BWP respectively.
 - Hazard a guess as to the number of pulas you would have to buy for the cost to be the same whether you use a credit card or cash. (To be continued later!)
- A leisure aircraft is owned by the Broadland Flying Group which has 10 members. The costs of keeping the aircraft comprise maintenance costs (including insurance) and the running cost which is mainly fuel. Average fuel consumption is 9 miles per gallon.

If the 10 members of the Group are to share costs equally, write down a formula to enable Gordon the treasurer to calculate the annual cost for each member if the aircraft travels n miles in a year, incurs maintenance costs of £ M and fuel costs £3 a gallon.



4. Consider the model for the clothing manufacturer

$$\text{Profit} = \text{price} \times \text{number sold} - (10,000 + 15 \times \text{number manufactured}).$$

Set up a spreadsheet to investigate the profit, when the selling price is £18, for a range of values for the number sold and the number manufactured. Comment on your results. Adapt your spreadsheet to investigate how many items must be manufactured in order to break even (make a profit of 0) assuming that the number sold is the same as the number manufactured.

6 **Working with symbols: adding, subtracting, multiplying, dividing****Test box 6**

Simplify the following by collecting like terms:

$$3pq + 2q + 2pq$$

Write the following expressions more succinctly:

$$3 \times (-2p) \times 2 \times p \quad 2 \cdot 3 \cdot 5$$

Solutions:

$$5pq + 2q$$

$$-12p^2 \quad 30$$

To construct models and solve equations we need to know how to manipulate symbols. This need not be a problem if we remember that the symbols merely *stand in for the numbers* and so can be treated in exactly the same way. In this section we recall the work of Sections 1–3, but apply it particularly to expressions containing symbols.

**Negative symbols**

The rules for adding, subtracting, multiplying and dividing numbers apply equally well to symbols. We can summarise these rules as

Adding and subtracting

SAME SIGNS

– (–a) or + (+a) gives +a

OPPOSITE SIGNS

+ (–a) or – (+a) gives –a

where a is any number, symbol or expression.

Multiplying and dividing

SAME SIGNS

multiplying $a \times b = ab$

$$(-a)(-b) = ab$$

dividing $a \div b = \frac{a}{b}$

and $(-a) \div (-b) = \frac{-a}{-b} = \frac{a}{b}$

OPPOSITE SIGNS

multiplying $a \times (-b) = -ab$

$$(-a) \times b = -ab$$

dividing $(-a) \div b = \frac{-a}{b} = -\frac{a}{b}$

$$a \div (-b) = \frac{a}{-b} = -\frac{a}{b}$$

Check these

$$2 + (-a) = 2 - a$$

$$3 - (-a) = 3 + a$$

$$2 \times (-a) = -2a$$

$$(-3) \times (-a) = 3a$$

$$(-b) \times c = -bc$$

$$(-c) \times (-a) = ac$$

$$\frac{p}{-q} = -\frac{p}{q} \quad \frac{a}{a} = 1 \quad \frac{-x}{-x} = 1 \quad \frac{-a}{a} = -1$$

Addition and subtraction: collecting 'like terms'

Because $a + a + 2a$ means one 'a' plus one 'a' plus two 'a's the terms can be collected together and written as $4a$. We can do this because each term is the same apart from the number – called the *coefficient*, which it is multiplied by. It is just like saying that 'one banana plus another banana plus another two bananas gives four bananas'.

Even when we have more complicated terms we can collect them together *as long as they are the same*. For instance

$$2pq + pq - 5pq$$

simplifies to $-2pq$, or

$$\frac{s}{2r} + 4\frac{s}{2r}$$

is equivalent to

$$5 \frac{s}{2r}$$

as each term is so many

$$\frac{s}{2r}s$$

Notice, however, that we *can't* collect together any terms in

$$3pq + p + q$$

as they are all multiples of different things – we can't add 3 bananas, an orange and a grapefruit!

Often just some of the terms in an expression can be collected together. For instance,

$$pq + 2p + 5pq$$

simplifies to $6pq + 2p$.

Check these

Where possible, simplify the following:

$$5rs - 3rs + rs$$

All the terms involve a number of rs , so the expression simplifies to $3rs$.

$$9pq + 4q - 5pq - q$$

The pq terms can be collected to give $9pq - 5pq = 4pq$, but the terms in q must be collected separately to give $4q - q = 3q$. So the expression simplifies to $4pq + 3q$.

$$3\frac{a}{2} - 2\frac{a}{2} + 5\frac{a}{2}$$

All the terms involve a number of $\frac{a}{2}s$, so the expression simplifies to $6\frac{a}{2}$. We will see later that this can be simplified further to $3a$.

$$2xz + 5zx - 3z$$

You should spot here that xz is the same as zx (as the order in which we multiply doesn't effect the result), so we can collect together $2xz + 5zx$ to give $7xz$. The whole expression simplifies to $7xz - 3z$.

In the last example it would have been easier to spot that $2xz$ and $5zx$ could be collected together if the xz and zx terms had been written in the same way. It is therefore a good idea to write letters that are multiplied together in alphabetical order. This is one of the suggestions below.

Some conventions for multiplying

1. We have already said that there is no need to include a multiplication sign '×' next to a bracket or next to a symbol. As a consequence, whenever letters or brackets appear

adjacent to each other a multiplication sign is implied. For instance $2ab(c + d)$ means $2 \times a \times b \times (c + d)$.

- We can't just drop the multiplication sign when multiplying two or more numbers as we wouldn't be able to tell when one number finished and the next one started. For example 357890 might mean 35×7890 or $3 \times 57 \times 890$ or many other possibilities. Instead we can shorten expressions by writing a slightly raised *dot* instead of the \times sign. So $357 \times 8 \times 90$ could be written $357 \cdot 8 \cdot 90$.
- When a number is multiplied by a letter it is conventional to write the number first so, for instance, we write $5n$ instead of $n5$.
- When a mixture of numbers and symbols are multiplied together, it is more succinct to multiply the numbers together so, for instance, instead of $4 \times p \times q \times 2$ we would write $8pq$.
- When several symbols are multiplied together it is usual to write them in alphabetical order. This means that 'like' terms can be spotted easily. So $cad \times 2$ would more usually be written $2acd$.
- When a number or symbol is multiplied by itself we say it is 'squared', and we usually write the number with a 2 superscript. For instance, 3^2 is called 'three squared' and means 3×3 . In a similar way $(ab)^2$ means $(ab) \times (ab)$.

Check this

Check that you understand what the following expressions mean

$$1. \quad 3p(2 + a) \quad \frac{100}{rc} \quad \frac{2 \cdot 3 \cdot 4}{ax} \quad \frac{100pq}{3 \cdot 2 \cdot 1} \quad \frac{a^2b}{c^2}$$

Write the following expressions more succinctly

$$2. \quad y \times z \times 2 \times a \quad 3 \times b \times a \times 2 \quad a \times 2b \times 5$$

Solutions:

- Writing out these expressions in long-hand using \times for multiplication gives: $3 \times p \times (2 + a)$, $100 \div (r \times c)$, $(2 \times 3 \times 4) \div (a \times x)$, $(100 \times p \times q) \div (3 \times 2 \times 1)$, $(a \times a \times b) \div (c \times c)$
- $2ayz$, $6ab$; remember that this is $a \times 2 \times b \times 5$, giving $10ab$.

Can I do this?

There are many more techniques for working with symbols and some will be considered in Chapter EM2. However, if you are in doubt as to whether two expressions are equivalent you can always evaluate both of them for some arbitrarily chosen numbers and see if the results are the same.

For instance, suppose you are unsure whether $p \times q$ is the same as $q \times p$. (We have picked an easy one to start with – we have already said that this is true.) If, for example, $p = 2$ and $q = 1$, then $p \times q = 2 \times 1 = 2$ and $q \times p = 1 \times 2 = 2$, so the expressions are equal for these values. Now try $p = -5$ and $q = 2$, putting a negative number to the test, and we have $p \times q = -5 \times 2 = -10$ and $q \times p = 2 \times -5 = -10$ which again are equal.

Be warned, however, that 'trying out' values like this does not constitute proof that the expressions are equal – you may just have been lucky and by chance selected values that



worked. If the two expressions are *not* equal for the values you have chosen, then this does, however, prove that they are *not* equivalent expressions.

For instance, is it true that

$$\frac{m+n}{n}$$

is the same as m ? Try some values for m and n to see whether this is true or not before you read on.

Check this

Is $\frac{m+n}{n}$ the same as m ?

Solution:

We'll choose $m = 100$, $n = 5$ first. When we evaluate the two expressions we obtain

$$\frac{m+n}{n} = \frac{105}{5} = 21$$

whereas $m = 100$. No, this does *not* work. We don't need to try any more values for m and n as we only need to find one counter-example to show that the two expressions are not equal.

WORK CARD 6

- Simplify, by collecting like terms where possible
 - $b + b + b + a + a$
 - $p + q + 2pq - q + 3qp$
 - $2p + 2q + p + pqr - 2pqr$
 - $2\frac{x}{y} + x - 2x + 3\frac{x}{y}$
- Write the following expressions more succinctly
 - $2 \times n \times 3 \times m$
 - $pq \times rs \times 3$
 - $a \cdot 2 \cdot 10 \cdot zb$
 - $(z \times y) \times (2 \times w)$
- You dimly recollect from your school career that it is valid to 'cancel down' and that

$$\frac{2a}{2b} \text{ is the same as } \frac{a}{b}.$$

Try out some numbers to see if this seems to be the case or not.

- Is it true that

$$\frac{1+2a}{2b} = \frac{1+a}{b} ?$$



Solutions:

1. a. $3b + 2a$ b. $p + 5pq$ c. $3p + 2q - pqr$ d. $5\frac{x}{y} - x$

2. a. $6mn$ b. $3pqrs$ c. $20abz$ d. $2wyz$

3. Try for instance $a = 5$ and $b = 10$, $2a = 10$ and $2b = 20$, so

$$\frac{2a}{2b} = \frac{10}{20} = 0.5 = \frac{a}{b}$$

so it works. Now try one or more negative numbers, say $a = -1$ and $b = 3$; then

$$\frac{2a}{2b} = \frac{-2}{6} = -\frac{1}{3} \quad \text{and} \quad \frac{a}{b} = \frac{-1}{3} = -\frac{1}{3}$$

so again it works. In fact it is true and we cover it in the next section!

4. No it is not true. Try for instance $a = 1$, $b = 2$; then

$$\frac{1+2a}{2b} = \frac{3}{4} \quad \text{whereas} \quad \frac{1+a}{b} = \frac{2}{2} = 1$$

which is clearly different.

1. Simplify by collecting like terms:

a. $3x - 2xy + 3y + 3xy$

b. $p - 2q - 2q^2 + q^2 - 3p^2$

c. $3 \cdot \frac{a}{2b} + a - 2\frac{a}{2b} + 3a$

2. Write down the following expressions more succinctly:

a. $\frac{f \times b \times 2 \cdot 3}{2 \times e \times d}$ b. $(c \times b) \times (a \times d) \times 3$

3. Investigate whether $a(b + d)$ is the same as $ab + d$.

Index

- μ *see* mu
 χ^2 *see* chi-squared distribution
 Σ *see* sigma
- A AND B event 344–9
A OR B event 344–9
ABC classification 753
absolute value 458
accrued interest 172–3
activity 726
 float 736
 preceding 730
addition of fractions 40–6, 54
additive model 627, 633,
 635–9
aggregate indices 316–318
alternative hypothesis 533–45
 contingency table 689
 linear regression model
 588–9
 one- and two-sided 545–6
 and rejection regions
 556–60
 testing for mean 549
AND events 344–9
AND probabilities 360–7
annual percentage rate 760
annuities 783
 discount rates 783
 present values 783
arrival rate 857
assignable causes, quality
 control 822
association, test of 689
averages, *see also* expected value
 run length 833
- back ordering *see* backlogged
 demands
backlogged demands 749–51,
 753
bar charts 271–4
base 2 logarithms 169
base 10 logarithms 169
base e logarithms 229–33
base periods 315
base values 314–15
base-weighted index 317
bases, logarithms 162–71
batch size 744–5, 749
binary data 673
 and categorical data 686–7
binding constraints 716, 717
binomial distribution 398–415
 cumulative probabilities
 405–6
 mean 408–9
 normal approximation to
 471–6
 plots 409–11
 probabilities 399–406
 probabilities, using a
 computer 408
 simulation 853–4
 summary 411–12
 tables of probabilities 406–8,
 862–6
 variance 408–9
 when n is large 471–8
brackets
 expanding 57–64
 introduction 11–12
 layers of 12
 multiplying out 57–64
 products of 61–4
break-even analysis 199–201
budget constraints, linear
 equations 147
business modelling 703–860
- calculators 14–19
 decimal places 16–17
 estimates 15–16
 ln key 165
 log key 165
 logarithms 165, 166
 operations, order of 14–15
 powers 88
 rounding 16–17
 scientific notation 17–18
 significant figures 16–17
 to calculate variance 306
calculus *see* differentiation
cancelling fractions 34–40
car-wash, simulation 841–4
categorical data 217, 672–98
 binary data 686–7
chi-squared test 681–98
 formulae 702

-value 686
 population proportion
 674–80

cells, spreadsheets 24–5
centre line
 control chart 822–34

-chart 832
 R-chart 831
centred moving averages 628,
 630–2
Chebysheff's result
 for data set 301–2
 for random variable 394
chi-squared distribution
 682–3, 873
chi-squared test 681–2
 and independence 691–2
 software 686, 692
classes 257
 frequency 319
 width 261–5, 318–20
co-ordinates, straight lines
 134–5
coefficient of determination
 578–80
coefficients 136
 of least squares line 577–82
 in linear equations 136–45
 in quadratic functions
 195–6
columns, spreadsheets 24
combinations 399–402
combining events, probability
 344–52
common denominator 42–6
common factors 38
comparing population means
 649–65
comparing two populations
 649–70
complementary events,
 probability 338–40
compound interest 759–65
 curve 212–13
 non-annual 760–5
 summary 764–5
computers 255, 263–75
computer software *see* Excel,
 SPSS statistical software
conditional probabilities
 352–60
 calculating 353–5
 dependent events 356–7,
 364–5

- independent events 356–60, 364–5
- terminal node 369
- on a Venn diagram 353–4
- confidence interval for proportion 674–6
- confidence intervals for
 - difference between two means 651–3, 659–60, 664–5
 - computer, on a 667–8
- confidence intervals for the mean 505–24
- on a computer 520–1
- interpretation 509–10
- large samples 520
- summary 520
- and two-sided tests 546–7
- variance known 505–16
- variance unknown 516–21
- confidence limits 508–9
- constants 22
- quadratic functions 194
- constraints, linear programming 707–13, 715–20
- contingency tables 281–2, 689
 - categorical data 689–702
 - chi-squared test statistic 689–92
 - expected frequencies 689–92
 - observed frequencies 689–92
- continuous compounding curve 213
- continuous distributions 438
- continuous random variables 377–8
 - expected value 436
 - exponential 440–1
 - probability distributions for 428–30
 - uniform 434–6
 - variance 436
- control charts
 - 2-sigma 824
 - 3-sigma 824
 - attributes 832–3
 - centre line 822–33
 - control limits 822–33
 - designing 833–4
 - for mean 820–37
 - for proportion 832
 - P*-charts 830–3
 - R*-charts 830–1
 - software 826–7
 - and testing 824
 - for variance 823, 832
- control limits 822–34
 - 2-sigma 824–5
 - 3-sigma 824–6, 832, 833
 - calculating 823–4
- controlling quality 820–37
- convergence, geometric series 778
- correlation 277, 563–610
 - coefficient 566–71
 - sample 566–71
- costs, inventory system 744–53
- courses of action, alternative 790–2, 794
- crashing, project planning 738
- critical activities, project planning 736–8
- critical paths, project planning 737–8
- crossing points, quadratic curve 195
- cube roots, powers 86–7
- cumulative binomial probabilities 405–8
 - tables 862–6
- cumulative normal probabilities 451–2
- cumulative standard normal probabilities, tables 867–8
- current-weighted price index 316
- curve sketching 217–18
- curved graphs 148–9
- curves
 - common functions 203–7
 - compound interest 213
 - continuous compounding 213
 - gradient of 221–8
 - inflexion point 248–50
 - maxima/minima 246–52
 - production 213–14
 - quadratic functions 192–208
 - sales 215
 - tangent of 222–4
 - turning/stationary points 242–53
- data
 - binary 673–4, 686–7
 - categorical 271–4, 674, 686–7
 - describing 256–326
 - distribution of 257
 - grouped in classes 318–22
 - numerical 271, 673
 - misrepresentation of 286–9
 - paired 277–85, 563–84, 663–5
 - pictures of 256–66
 - qualitative 271
 - quantitative 271, 673
 - sequential 271, 278
 - summarising 291–326
- data set *see* data
- decimal places, calculator 16
- decision analysis 790
- decision making 789–819
 - courses of action 790
 - criteria 797–803
 - objective 790–2
- decision trees 804–18
 - branches 805
 - drawing 805–12
 - nodes 805
- decision variables, linear programming 707–8, 711–20
- demand
 - Economic Order Quantity 744–53
 - elasticity of 237–9
 - inelastic 239
- denominator 35
- density functions 427–38
- dependent events 356, 364–5
- joint probability of 363
- dependent variables, regression 577
- derivatives 225–54
 - interpretation 234–40
 - notation 226–7
 - second 245–52
 - see also* differentiation
- describing data 256–326
- descriptive statistics 481–2
- deterministic models 841
- differentiation 221, 225–54
 - of a difference 230–1, 233
 - and gradients 225–6
 - of a logarithm 229–30, 232
 - of a multiple 231–2
 - notation 226–7
 - summary 232–3
 - of a sum 230–1, 233
- discount rate 767–72
- discrete random variables 377–9, 427–8
 - probability distribution 430
- discrete uniform distribution 854
- distributions *see also* probability distributions
 - of a set of data 257
 - of a population 490–1
 - sampling distributions 495
 - skewed 295–6
- division
 - by zero 8, 106

- of fractions 46, 50–3
- of powers 75–7
- dummy
 - activity 728–35
 - logical 728–9
 - uniqueness 730
- earliest event times 733–6
- Economic Order Quantity (EOQ) 744–55
 - backlogged demand/back ordering 749–51, 753
 - batch size 744, 749
 - demand 744–52
 - formula 745, 751
 - lead time 749
 - reorder level 749
 - shortages/stock out 749
 - variability coefficient 752–3
- effective interest rate 760–5
- elasticity of demand 237–9
- EOQ *see* Economic Order Quantity
- equations
 - equivalent 101–4
 - formulation of 94, 97–8, 108–11
 - guessing solutions 97–101
 - linear 135–8, 147–56
 - for problem solving 93–132
 - rearranging 111–14
 - solution of 97–100, 108–11
 - straight lines 133–7, 143–4, 151
 - subject of 112–15
 - substitution 95–6, 111, 118–20
 - transposing 112–14
- equivalent fractions 34–40
- error sum of squares, in regression 577–82
- errors
 - forecasting 619–26
 - in hypothesis testing 534–7
 - Type I error 535–7
 - Type II error 535–7
- estimates *see also* estimation
 - rough 15–16
- estimation 483–524, 699
 - of difference between two means 651–3, 659–60, 664–5
 - of linear regression model 586–7
 - of multiple linear regression model 603
 - of population mean and variance 491–2
 - of process mean and variance 825–6
 - of trend 632, 633–7
- event times
 - project planning 733–6
 - simulation 842–4, 849, 856
- events
 - A AND B 344–9
 - A OR B 344–9
 - combining 344–52
 - complementary 338–40
 - dependent 356, 363–5
 - independent 356–65
 - mutually exclusive 348–9, 357
 - Poisson distribution 415–17
- Excel software 255
- bar charts 274
- basic statistics 311
- binomial probability 408
- chi-squared test statistic 692
- comparing population mean 668
- exponential forecasting 623
- exponential probability 447
- frequency distribution 265
- generating probability distributions 855–6
- generating random numbers 848
- histogram 265
- internal rate of return 771–2
- least squares line 581
- linear programming 715–17, 718–20
- linear regression model 592–4
- moving averages 632
- net present values 771–2
- normal probability 468
- payments, series of 784–5
- pie charts 274
- Poisson probability 421
- sample correlation 571
- sampling 488
- scatter plots 278
- time series plots 278
- expanding brackets 57–64
- expectation *see* expected value
- expected frequencies
 - categorical data 680–8
 - contingency tables 689–92
- expected value 384–7
 - compared with \bar{x} 387
 - of a continuous distribution 436–8
 - of difference between means 650–1
 - of a function of X 388–9
 - as long run average 384–5
 - of sample mean 496–7
 - of X^2 387–8
- expected value criterion 798–801, 812–14
- exponential distribution
 - 440–8
 - mean 443
 - and Poisson distribution 441–2
 - probabilities 444–7
 - probabilities using a computer 447
 - simulation 854
 - variance 443
- exponential smoothing 619
- F test 606
- factorial 401
- factorising 64–9
 - benefits of 65
 - quadratic equations, to solve 179
 - quadratic expressions 66–9
- factors 64
 - common 38
- feasible regions, linear programming 710–15, 720
- final nodes, project planning 728
- fixed order quantity 117
- fixed-increment time-advance mechanisms 849
- floats, project planning 736
- forecasting
 - additive model 644–5
 - errors 619–23
 - estimating trends 633–4, 635–6, 639
 - exponential 617–24
 - four period moving average 628–31
 - mean absolute error 619–26, 633–5, 641–2, 646
 - mean squared error 619–26
 - moving average 628–33
 - multiplicative model 645
 - notation 616–17
 - seasonal effects 627–44
 - seasonal indices 633–7, 639
 - seasonally adjusting series 637–8
 - smoothing constant 618–19, 622–3
 - trend and seasonal effects 627–44
- formulae 21

- categorical data 701–2
- combination 399–400, 403–5
- compound interest 759, 763
- correlation 700
- Economic Order Quantity 745, 751
- expected value 385
- forecasting 701
- linear regression model 586
- normal distribution 449
- Poisson distribution 417
- present values 767, 772
- probability summary 479–80
- quadratic equations 176–9, 183
- quadratic functions 175
- regression 700–1
- sample correlation 571
- sample mean 306
- statistics summary 699–702
- testing hypotheses 700
- variance 306
- formulating
 - equations 94, 97–8, 108–10
 - inequalities 123–9
 - linear programming models 707–8, 717–20
- four period moving averages 628–31
- fractional powers 85–90
- fractions 34–56
 - addition of 40–5, 54
 - cancelling 38–9, 48–50, 52
 - common denominator 42–5
 - division of 46, 50–3
 - equivalent 34–40
 - lowest common denominator of 42–5
 - lowest terms 37–40
 - multiplication of 46–51, 52, 54
 - operations, order of 54
 - quotients 35
 - reciprocals 51
 - simplest terms 37–40
 - subtraction of 40–5, 54
- frequencies 258, 272, 319
 - on a computer 263–4
 - expected 681
 - observed 680
 - relative 258
- frequency distribution 265
- functions 189–91
 - curved L-shaped 206
 - demand 241
 - expected value of 388–9
 - inverse 165–6
 - linear 134–7, 191
 - marginal product of labour 236–7
 - marginal revenue 235–6
 - maxima/minima of 221
 - non-linear 189, 196–223
 - objective 707–13, 715, 720–1
 - turning/stationary points 250
 - see also* quadratic functions
- Gantt charts 737–8
- geometric ratios 776, 779
- geometric series
 - present values 781–2
 - sums accrued 776–9
- global maxima/minima 242–5
- gradient
 - of a curve 221–8
 - and derivatives 226
 - and differentiation 227
 - negative 143
 - positive 143
 - of straight line 141–44
 - and tangents 223–4
- graph sketching 134–8
 - functions 192–224
 - quadratic functions 196–9
 - straight lines 137–9
- grouped data 318–24
- histograms 257–65
 - class width 261–3
 - number of classes 263
 - and probability density functions 430–2
 - and relative frequencies 257–65, 271–5
 - skewed distributions 260, 265, 269, 295–7
 - software 263–5
 - symmetric distributions 260, 295–7
- holding costs 744–7, 750–1
- hypotheses 527–8, 533
- hypothesis testing *see* testing
- imaginary solution to equation 177
- independence, test of 691–2
- independent events 356
 - joint probabilities for 361–5
 - and mutually exclusive events 357
- independent variables, regression 577
- index numbers 314–17
- indices
 - aggregate 316–17
 - constructing 314–15
 - powers 70–91
- inelastic demand 239
- inequalities 123–32
 - applications of 124–5
 - rearranging 125–8, 171–2
 - signs 123
- inferential statistics 481–3
- inflexion point 248–50
- initial conditions, simulation 843
- inputs, simulation 841–47
- instalments, loan repayment 781–2
- integer programming 713
- inter-arrival times 842–4, 856
- inter-quartile range 309–10
- intercepts, with y axis 140–1, 144
- interest rates
 - annual percentage rate 760
 - discounting 767
 - internal rate of return 770–2
 - net present value 769–72
 - nominal 760
 - present value 767–74
 - principal 758
- internal rate of return (IRR) 770–2
 - for series of payments 784
 - software 771–2
- interpolation 309
- interval estimates 505–15
 - see also* confidence intervals
- inventory control 742–56
 - ABC classification 753
 - batch size 749
 - costs 744–53
 - holding costs 744–7, 750–1
 - lead time 749
 - ordering costs 744–7
 - set-up costs 744–7, 750
 - shortage costs 744, 749–51
 - unit costs 744–53
 - see also* Economic Order Quantity
- inverse functions 165–6
- inverse normal probabilities 455
- IRR *see* internal rate of return
- isolating trends
 - centred moving average 628, 630–2
 - regression 632–3
- iterative method 715

- joint probability 344
 - calculating 360–7
 - of dependent events 363–4
 - for independent events 361–4
- Laspeyre price index 316
- latest event times 733–7
- lead times, Economic Order Quantity 749
- leaf units 267–70
- least squares line 574–582
 - coefficient of determination 577–82
 - dependent variable 577
 - error sum of squares 577–82
 - errors 575
 - estimates 575
 - fitting 575
 - prediction 576
 - regression sum of squares 577–82
 - software 580–1
 - summary 582
 - total sum of squares 577–82
- linear functions 135–6
- linear equations
 - budget constraint 147
 - coefficients 136–9, 147–8
 - modelling 146–9
 - pairs of 150–7, 711
 - simultaneous 150–1, 156
 - solving pairs of 151–4, 156
 - and straight lines 135–7, 151–3
 - supply and demand models 146
 - types of solution 154–6
- linear programming models
 - binding constraint 716, 718
 - constraints 707–13, 716–23
 - decision variables 707–8, 712–20
 - feasible region 709–15, 720
 - graphical solution of 708–11
 - iterative method 715–16
 - limitations of 712–13
 - non-binding constraint 716, 718
 - non-negativity constraint 708, 718
 - objective function 707–8, 710–12, 715, 720
 - optimal solution 711–12, 715, 719–20
 - optimisation 711–12
 - problem formulation 707–8, 717–19
 - sensitivity analysis 719–20
- simplex method 715
- simultaneous equations 711
- software 715–16, 718–19
- unbounded solution 712
- unique optimal solution 712
- linear regression model 585–606
 - assumptions 585–6
 - and correlation 580
 - estimating 586–7
 - F test 605–6
 - interpretation 588
 - multiple 602–6
 - prediction 588
 - simple 585–601
 - software 592–4, 603–6
 - summary 594–5
 - testing 589–92
 - for trends 632–3
- In key 165
- loan repayments 781–2
- local maxima/minima 242–5
- log key 165
- logarithms 161–74, 182–3
 - bases 162–71
 - on calculator 165, 166
 - differentiation of 229–30, 232
 - evaluating 163–4, 166
 - inequalities 171–2
 - inverse function 165–6
 - manipulation of 164–5, 167
 - and powers 165–6
 - solving equations 168–72
- logical dummy activities, project planning 728–9
- logs *see* logarithms
- lower control limits
 - control chart 822–33
 - p*-chart 832–3
 - R*-chart 830–1
- lower quartiles 309–10
- lowest common denominator 42–5
- lowest terms, fractions 37–40
- MAE *see* mean absolute errors
- manipulating logarithms 164–7
- marginal functions 234–7
- marginal product of labour function 236–7
- marginal revenue function 235
- market equilibrium 156–7
- matched data *see* paired data
- maxima
 - global 242–4
 - local 242–4
- second derivatives 245–6
- maximax criterion 797
- maximin criterion 797–8
- maximising 242–53
- mean
 - on a computer 311
 - of data in classes 320–1
 - of population 491
 - of random variable 385–7
 - of sample 492
 - of a set of data 293
 - see also* expected value
- mean absolute error (MAE) 578–9, 619–26, 633–5, 646
- mean squared error (MSE) 619–26
- median 293–8, 320
- minima
 - global 242–4
 - local 242–4
 - second derivatives 245–6
- minimax regret criterion 798
- minimising 242–53
- MINITAB statistical software 255
- minus numbers *see* negative numbers
- mode
 - of data 295–6, 298, 320
 - spreadsheets 24–5
- moving average 638–31
 - centred 628–32
 - forecasting 629–32
 - software 632–3
- MSE *see* mean squared error
- mu (μ) 385
 - and sample mean 387
- multiple linear regression 602–6
 - assumptions 602
 - interpretation 604
 - test statistic 606
- multiples, differentiation of 231–3
- multiplication
 - brackets 57–64
 - by zero 106
 - fractions 46–50, 51–2, 55
 - powers 72–4, 77
- multiplicative model 627, 633–5, 639, 645
- mutually exclusive events 348–9, 357
 - Venn diagram 349
- negative
 - gradient 143
 - powers 72, 77

- negative numbers
 - addition 4–6
 - division 7–8
 - multiplication 6–7
 - sign conventions 7
 - subtraction 4–6
- net present value (NPV)
 - 769–72
 - of series of payments 784
 - software 771–2
- networks, project planning
 - 727–41
- next-event time-advance
 - mechanisms, simulation 849
- nodes
 - project planning 727–41
 - tree diagram 368–9
- nominal rates of interest 759–60
- non-annual compound interest
 - effective rate 759–62
 - investment value 762–4
- non-binding constraints
 - 716–18
- non-convergence, geometric series 778
- non-critical activities 737–8
- non-linear functions 189, 197–223
- non-negativity constraints
 - 708, 718
- normal distribution 449–71
 - approximation to binomial distribution 471–6
 - formula 449
 - mean 450
 - notation 449–50
 - probabilities 451–68
 - using a computer 468
 - simulation 853–4
 - standardising 460–1
 - standard normal 450–60
 - tables 452, 869–70
 - variance 449–50
- NPV *see* net present values
- null hypothesis 533–7
 - contingency tables 689–92
 - linear regression model 589
 - and rejection regions 555–60
 - testing for mean 539–40, 654
 - testing for proportions 676, 680
- numerator 35
- numerical data 271
- objective function 707, 711, 716
- observed frequencies
 - categorical data 680–8
 - contingency table 689–92
- one-sided alternative hypothesis
 - 545–6
- operations, order of 10–15, 54
- opportunity loss 798
- optimal solution 712–13, 715, 720
- optimisation, linear
 - programming 712–13
- OR events 344–9
- ordering costs 743–7
- outputs, simulation 842–4, 856
- p*-charts 830–3
 - centre line 832
 - lower control limit 832–3
 - software 833–4
 - upper control limit 832–3
- p*-values 533–4
- Paasche index 316
- paired data 277–8, 281–5, 564–601
 - displaying 277–8
- paired samples 663–70
 - confidence interval 664–5
 - testing 663–5, 667
- pairs of linear equations
 - 150–7
- parameters, linear regression
 - model 586, 603
- patterns, control charts 823
- payments, series of 775–88
 - internal rate of return 784
 - present value of 781–4
 - software 784–5
- payoffs 790–2, 794, 809
- payoff tables 793–6, 800
- pdfs *see* probability density functions
- percentage points
 - chi-squared distribution 683, 873
 - normal distribution 465–8
 - standard normal distribution 454–5, 869–70
 - t* distribution 517–18, 872
- percentages 55–7
 - for confidence interval 511–14
 - reporting 282–3
- pictorial representations
 - 257–66, 319–20
- pie charts 274
 - software 274
- planning projects 726–41
- plots
 - scatter 277–8, 565
 - time series 612–16
- Poisson distribution
 - approximation to binomial 418
 - and exponential distribution 441–2
 - mean 421
 - probabilities 417
 - on a computer 421
 - rare events 415–16
 - simulation 854
 - summary 421
 - variance 421
- pooled variance estimator
 - 657–8
- population 292
 - distribution 490–1
 - mean 491
 - proportion 673–4
 - variance 491
- population mean 491, 497
 - confidence interval 505–24
 - testing 539–62
- population proportion 673–4
 - confidence interval 674–6
 - testing 676–8
- population variance 491, 502
- positive
 - gradient 143
 - numbers 4, 6
- powers 70–91
 - applications 74, 88–9
 - on a calculator 88
 - cube roots 86
 - division of 75–6
 - fractional 85–91
 - indices 161–2
 - multiplication of 72–4, 77
 - negative 72, 77
 - of a power 74–5, 77
 - of products 80–5
 - of quotients 80–5
 - square roots 86
- precedence table 727–30
- predictions
 - least squares line 576
 - linear regression model 588
- preliminary samples 825
- present values 767–9, 772
 - annuity 783
 - of series of payments 775–6, 781–4
- price elasticity 237–9
- principal 758
- probability
 - of AND event 344–9

- complementary events 338–40
- cumulative 405–6
- inverse normal 455
- of OR event 344–9
- subjective 332–3
- and testing 529–31
- probability density functions (pdfs) 427–56
 - for discrete random variables 430
 - drawing 432–4
 - exponential distribution 440–1
 - and histograms 430–2
 - normal distribution 449–6, 453
 - uniform distribution 434–6
- probability distributions 378–9, 673
 - binomial 380, 398–415
 - chi-squared 683
 - of continuous random variables 428
 - and distributions of data 381
 - exponential 440–7
 - normal 449–71
 - Poisson 380, 415–21
 - simulation 846, 853–5
 - uniform 434
- process mean 822–6
- process variance 823–6
- production
 - costs 146–7
 - curves 213–14
- products
 - of brackets 61–3
 - of powers 80–5
- project planning 725–41
 - activity float 736
 - crashing 738
 - critical activity 736–8
 - critical path 737–8
 - dummy activity 728–30
 - event times 733–7
 - floats 736
 - Gantt chart 737–8
 - logical dummy activity 728–9
 - networks 727–41
 - node 727–41
 - non-critical activities 737–8
 - precedence table 727–30
 - scheduling activities 733–8
 - uniqueness dummies 730
- proportions 672–8
 - confidence interval 674–5
 - testing 676–8
- quadratic curves 192–208
- quadratic equations 161, 174–87
 - definition 174–5
 - factorising 179
 - imaginary solutions 177
 - no solution 177
 - real solution 177
 - solving 176–86
- quadratic expressions 66–9
- quadratic functions 192
 - break-even analysis 199–201
 - plotting 194–9
 - sketching 194–9
- qualitative data 271
- quality control 500–1, 820–37
 - assignable causes 822
 - average run lengths 833–4
 - p -chart 830, 832–4
 - process mean 825–7
 - process variance 825–7
 - R -chart 830–4
 - and statistical testing 824–5
- quality management 821
- quantitative data 271
- quantity discounts 746
- quartiles 309–10, 320
- queues, simulation 856–7
 - assumptions 856–8
 - single-server 856–7
- queuing problems 856–7
- queuing theory 856–7
- quotients 12
- powers of 80–4
- R^2 *see* coefficient of determination
- R -charts
 - centre line 830
 - lower control limits 830–2
 - upper control limits 830–2
- random inputs, simulation 842–4
 - generating 846–8
- random number tables 487, 846–8, 871
- random variables 375–88
 - Chebysheff's result 394
 - continuous 377–8
 - discrete 377–8, 427–8
 - expectation of 384–91
 - expected value of 385–91
 - mean of 385–9
 - and probability distributions 379–84, 430
 - standard deviation of 392, 395
 - variance of 391–8
- range, data sets 299
- rates of change 221–54
- rearranging equations 111–14
 - inequalities 125–8, 171
 - logarithms 171–2
- reciprocals 50
- regression *see* linear regression
- regression sum of squares 578–82
- rejection regions 555–60, 667
 - advantages 559–60
 - and p -values 555
 - testing for mean 555–60
- relative frequencies data sets 257–65, 271–4
- relative frequency approach 331, 353
- reorder level, Economic Order Quantity 749
- repaying loans 781–2
- residuals, standardised 684
 - analysis of 592
- rewards 790–2
- roots, powers 86–7
- rounding numbers 16–17
- 2-sigma control limits 824
- 3-sigma control limits 824–6, 832–4
- sample correlation, r 566–71
 - on a computer 570–1
 - and R^2 580–2
 - summary 571
- sample mean, \bar{x} 293, 295–6, 491
 - distribution of 497–8
 - mean 497
 - variance 497
- samples 292, 484–5
 - selecting 486–9
 - simple random 485–6
- sample variance, s^2 298–306, 491
 - distribution 502
- sampling
 - bad 488–9
 - distribution 495
 - fluctuation 492–3
 - selecting a simple random sample 486–8
- scatter plots 277–8, 565–6
 - software 278
- scheduling projects 725–41
- scientific notation 17–18

- seasonal effects 612–13, 627–44
- seasonal indices 633–9
- seasonal adjustment 637–8
- second derivatives 245–52
 - evaluation 245–6
 - maxima/minima 246–7
 - notation 245
- sensitivity analysis 719–20
- service rate 857
- set-up cost 743–7, 750
- shortage cost 743, 749–51
- sigma (Σ) 304–6
- signs
 - inequalities 123–4
 - negative numbers 6–8
 - symbols 27–8
- significance level
 - hypothesis testing 534–7
 - quality control 824–5
- significance probabilities *see p-values*
- significant figures 16–17
- simple interest 758–9, 764–5
- simple random sampling 485–90
- simplest terms 37–40
- simplex method 715
- simplifying expressions 33–91
- simulation 839–59
 - event times 842–4, 849, 856
 - generating random inputs 846–53
 - initial conditions 843
 - inter-arrival times 842–3, 856
 - next-event time mechanism 849
 - outputs 842–4, 856
 - probability distributions 853–8
 - and queuing problems 857
 - random inputs 841–2
 - single-server queue 857
 - summary statistics 844–5
 - time mechanisms 849–51
 - transient period 852
 - warmup period 852
- simultaneous linear equations 150, 156, 711
- single variable graphs 191
- single-server queues 857
- skewed
 - binomial distributions 409–10
 - histograms 260, 265, 269, 295–7
- slack, linear programming 716–19
- slope, straight line 141–4
- smoothing constants 618–23
- software *see* Excel; SPSS
 - statistical software
- solving equations 98–111
 - guessing 97
 - guidelines 104
 - pairs of linear equations 151–6
 - quadratic 176–86, 195–6
 - unknown powers 168–72
- spread, data sets 298–304
- spreadsheets
 - cells 24–5
 - columns 24
 - modes 24
 - rows 24
 - symbolic expression 24
 - values mode 24
 - to calculate variance 306
- SPSS statistical software
 - binomial distribution 408
 - bar charts 273–4
 - categorical data 273–4
 - chi-squared tests 686, 692
 - confidence intervals 520–1, 667
 - contingency tables 284
 - control charts 826
 - descriptive statistics 311, 650
 - exponential probabilities 447
 - exponential smoothing 623
 - histograms 264–5
 - least squares lines 581
 - linear regression 593–4
 - moving averages 632
 - multiple linear regression 603–6
 - normal probabilities 468
 - paired samples 667
 - pie charts 274
 - Poisson distribution 421
 - random numbers 486–7, 855
 - sample correlation 570–1
 - sampling 488
 - scatter plots 277–8
 - stem and leaf diagrams 269
 - time series plots 278
 - normal random variable 493
 - testing means 547–9
 - t intervals 520–1
- square roots 86
- standard deviation for control
 - charts 823–4, 832
 - of sample 300
 - of random variable 392, 395
- standard error 497
- standard normal distribution
 - 450–60
 - percentage points 869–70
 - probability density function 453–4
 - tables 452–4, 867–8
- standardised residuals 684
- start nodes 727
- stationary points
 - classification of 246–7, 248–50
 - curve sketching 247–8
 - finding 242–53
- statistical process control 820–837
- statistical tables
 - cumulative binomial probability 862–6
 - cumulative standard normal probability 867–8
 - percentage points chi-squared distribution 873
 - percentage points standard normal distribution 869–70
 - percentage points t distribution 872
 - random numbers 871
- statistical testing *see* testing
- statistics 481–702
 - descriptive 481
 - formulae 699–702
 - inferential 481–4
- stem and leaf diagrams 267–71
- stem units 267–70
- stochastic simulation 841
- stock exchange index 316
- stock levels, evolution of 750
- stock out 749
- straight lines
 - equation 134–7, 143–4, 151
 - fitting to data 574–82
 - functions 133–7
 - gradient 141–4
 - graphs of 134–5, 139
 - linear equations 135–7, 151
 - sketching 137–8
 - slope of 141–4
- subjective probabilities 332–3
- subject of equation 112–16
- substitution
 - equations 95–7, 111
 - expressions 118–22
 - symbols 21–2
- subtraction of fractions 40–6, 54

- summarising data 291–326
- summary statistics simulation 844–5
- summation sign *see* sigma
- sum accrued 775–80
- sums, differentiation of 230–1, 233
- supply and demand models
 - linear equations 146
 - quadratic equations 176
- symbolic expressions, spreadsheets 24
- symbols 20–1
 - addition 27–9
 - collecting like terms 28–9
 - division 28
 - equivalent expressions 30–1
 - fractions 38–9
 - multiplication conventions 28–30
 - negative 27–8
 - subtraction 27–9
- symmetric distribution 260, 295–8
- symmetry
 - binomial distribution 409–10
 - quadratic curves 193–201, 204
- t* distribution 516–19
- t* intervals 516–21
- t* ratios, linear regression 589–92
- tables *see* statistical tables
- tangents
 - to curve 223–4
 - and gradients 223–4
- terminal nodes 368–9
- testing 525–62, 668
 - chi-squared test 681, 686–7
 - on a computer 547, 667–8
- difference between means 649, 653–8, 664–8
- the idea of 526–7
- in linear regression 588–91, 604–6
- for mean 541–7
- for proportion 676–8
- structure of a test 527–8, 533–7
- test statistic 533, 540–1
- time series 612
 - models 627–8
 - plots 612–15
- total sum of squares 578–82
- transient period 852
- transposing equations 111–14
- tree diagrams
 - construction 368
 - nodes 368–9
- trends 612–16, 627–44
 - isolation of 627, 630–3
- turning points *see* stationary points
- two-sided alternative hypothesis 545–6, 655
- two-sided tests 546–7, 655
- two-tailed tests 546–7
- Type I error, hypothesis testing 535–7
- Type II error, hypothesis testing 535–7
- U-shaped functions 194–8, 204
- unbounded solution, linear programming 712
- uncontrollable events 790–2, 794, 807–8
- uniform distribution 434–6, 854
 - simulation 854
- unique optimal solutions 712–13
- uniqueness dummies 730
- unit cost, inventory control 744–53
- upper control limits
 - control chart 822–33
 - p*-chart 832–3
 - R*-chart 830–2
- upper quartiles 309–10
- variability coefficient, inventory control 752–3
- variables 22
 - dependent 577
 - independent 577
 - see also* random variables
- variance
 - calculating 306
 - and Chebysheff 394
 - of a continuous distribution 436–8
 - of a random variable 391–4
 - of S^2 501–2
 - of a sample 298–300, 304–6
 - of sample mean 496–7
 - of a set of data 298–300, 304–6
- Venn diagrams 337–8
 - A AND B 345, 347
 - A OR B 346–7
 - conditional probability 353–5
 - mutually exclusive events 349
- warmup period, simulation 852
- width, class 261–5, 318–20
- z* intervals 505–16